



## Geochemistry and Quality Characterization of Efon Psammite Ridge Spring Water, Southwestern Nigeria

Nwankwoala HO<sup>1</sup>, Nwaogu C<sup>2</sup>, Bolaji TA<sup>1</sup>, Uzoegbu MU<sup>1</sup>, Abrakasa S<sup>3</sup> and Amadi AN<sup>3</sup>

<sup>1</sup>Department of Geology, College of Natural and Applied Sciences, University of Port Harcourt, Nigeria

<sup>2</sup>Department of Environmental Geosciences, Czech University of Life Sciences, Kamycka 1176, 1652, Praque 6, Czech Republic

<sup>3</sup>Department of Geology, Federal University of Technology, Minna, Nigeria

### Abstract

This study evaluates the geochemistry and quality characterization in the Efon Psammite Spring, Southwestern Nigeria. Fifteen (15) water samples were collected from the study area at regular intervals and samples were analyzed in order to determine their quality characteristics. Except for pH values that is mildly acidic to slightly alkaline in some locations, the physico-chemical properties are below the WHO recommended standards for drinking water. The low values of the conductivity are mainly attributed to geochemical processes prevailing in the area. The mean concentration of the cations follows the order:  $\text{Na}^+ > \text{Ca}^{2+} > \text{K}^+ > \text{Mg}^{2+}$  while for anions,  $\text{HCO}_3^- > \text{Cl}^- > \text{PO}_4^{3-} > \text{SO}_4^{2-} > \text{NO}_3^-$ . The water is mildly acidic to alkaline due to dissociation of bicarbonate with the following water types:  $\text{Na-SO}_4\text{-Cl}$ ,  $\text{Na-HCO}_3$ ,  $\text{Ca-Na-SO}_4$  and  $\text{Ca-Mg-HCO}_3\text{-SO}_4$  which are a reflection of geology and climate of the area. The mildly acidic to alkaline indices revealed that the spring water has undergone ion exchange between  $\text{Na}+\text{K}$  ions in the water with  $\text{Ca}$  and  $\text{Mg}$  of soil during the residence time of the water. The springs are being recharged from recent precipitation that has low water-rock interactions and low residence time within the aquiferous zones. The hydrochemical trend signifies low mineralized water with low water-rock interactions and residence time. Based on these water types and the presence of  $\text{Na}^+$ , the concentrations of cations are geogenic in origin and might have come from the interaction of water and the rock or introduced from weathering of rocks into the spring water. It is recommended that effective development of springs should involve thorough examination of their seasonal discharges, including during the summer. In case water supply exceeds use, the surplus waters may be stored for future use in horticulture and to irrigate crop land. Moreso since springs are yet to receive substantial attention, utmost care must be taken to protect the Efon Psammite spring from contamination.

**Keywords:** Water quality; Spring water; Geochemistry; Efon; Nigeria

### Introduction

Springs are natural outlets through which groundwater emerges at the ground surface as concentrated discharge from an aquifer are one of the most conspicuous forms of natural return of groundwater to the surface [1]. Conditions necessary to produce springs are many and are related to different combinations of geologic, hydrologic, hydraulic, pedological, climatic and even biologic controls [2]. Springs while flowing downhill from their origins, have formed their own narrow riparian zones on either side of their flow-paths. These riparian zones usually have heavy vegetation growth. Undoubtedly, these plant growths and the seeping spring waters augment the groundwater recharge of the blower aquifers to support the life of the existing springs issuing from these aquifers. Freshwater from the spring could be directly discharged onto the ground surface, directly into the beds of rivers or streams or into the ocean below sea level. Springs are used both for drinking and irrigation purposes by the local inhabitants both at higher elevations and in foothill zones (Figure 1). Spring water was associated in the public mind with exceptional quality and even considered holy in some places. Bottling of spring water has become a blooming business across the world [3]. Apart from tapping springs for drinking/irrigation purposes, they also sustain thousands of other life forms vital to a balanced ecosystem [1]. The importance of springs have gone beyond just being sources of domestic and municipal water supply but also sources for foreign exchange earnings as they serve as places for tourist attraction and industrial establishment where safe drinking water could be bottled [4]. Water chemistry to a large extent, is influenced by elemental distribution often determined by the lithologic effects, climate, groundwater flow and anthropogenic activities [5,6].

The sources of water for any specific purpose are not as important as the suitability of the water for the desired purpose as the exposed springs can be contaminated through anthropogenic activities. With

increasing human population, industrialization, urbanization and the consequent increase in demand for water for domestic and industrial uses, the increase in the implication of polluted water on man and the environment have been on the increase and have been severally studied [7,8]. Assessment of Efon Psammite spring water is therefore necessary as increased knowledge of processes that controls chemical compositions of the spring water can improve the understanding of its usability status apart from guide against outbreak of water borne diseases. This study is therefore aimed at evaluating the quality characteristics as well as examining the possible use of Efon Psammite ridge spring as a potable source of water (Figure 2).

### The Study Area, Geology/Hydrogeology

The study area (Ekiti Basement Complex Area) is within the Precambrian-Lower Proterozoic Basement Complex terrain of southwestern Nigeria [9,10]. It is located on coordinates: Latitudes  $7^{\circ}38.678'N$  and  $7^{\circ}49.914'N$ , Longitudes  $4^{\circ}54.658'E$  and  $4^{\circ}57.478'E$ . Highest elevation is about 523 m above mean sea level. Efon Psammite is an extensive quartzite ridge that cut across Ido, Ajinare, Ita-Ido,

**\*Corresponding author:** Nwankwoala HO, Department of Geology, College of Natural and Applied Sciences, University of Port Harcourt, Nigeria, Tel: +234-8036723009; E-mail: [nwankwoala\\_ho@yahoo.com](mailto:nwankwoala_ho@yahoo.com)

**Received** October 15, 2015; **Accepted** October 26, 2015; **Published** November 01, 2015

**Citation:** Nwankwoala HO, Nwaogu C, Bolaji TA, Uzoegbu MU, Abrakasa S, et al., (2015) Geochemistry and Quality Characterization of Efon Psammite Ridge Spring Water, Southwestern Nigeria. J Geol Geophys 4: 229. doi:[10.4172/jgg.1000229](https://doi.org/10.4172/jgg.1000229)

**Copyright:** © 2015 Nwankwoala HO, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1: Map of Ekiti State showing the study area-Effon.




Figure 2: Geologic Map of Ekiti-State, Nigeria.

Itawure, and Effon Alaaye.

The principal rock units in the area include Precambrian migmatite-gneiss-schist complex constituting over 60% of the sampling localities into which other rocks intruded during the Pan-African orogeny. Among the rocks that intruded into the migmatite-gneiss-schist complex were the charnockite, fine grained granite and porphyritic granite. All the rocks which have previously been described as Younger metasediments belong in this group e.g., Effon Psammite Formation [9]. The Effon Psammite Formation is a belt of quartzites, quartz schists and granulites which occurs largely east of Ilesha and runs for nearly 180 km in general NNE-SSW direction. The topography of the area is the undulating type, dotted with different outcrops in several places. Most of the rocks in the area are well exposed and are as high as 250 m above sea level and there is prevalence of erosion gullies along hill slopes and valleys.

The drainage system over the area consisting of the basement rocks are usually marked with the proliferation of many smaller river channels. The channels of these smaller streams are dry for many months especially from November to May. There is a major river in the study area called River Oyi and Owawa in Aiyegunle village. The drainage pattern in the south-eastern part of the area where topography is dominated by series of ridges is the trellis type which suggests that the drainage here is structurally controlled whereas, the drainage pattern in other parts of the study area is dendritic, due to homogeneity of the rocks and absence of structures. The Effon Psammite is a NNE-SSW

trending ridge highly foliated, jointed, schistose ridge often showing interesting sedimentary structures. The intense deformation structures of these rocks permit adequate aquifer properties needed to generate the springs. Consequently, the quartzite belt stretching NNE-SSW is considered as good aquifer. The surrounding schistose units occur as low yielding aquicludes. Ekiti State has so many springs, amongst which are: Ikogosi warm spring in Ikogosi-Ekiti, Arinta spring in Ipole-Ekiti and Arioye and Afeni springs in Efon-Alaye-Ekiti. The springs have history of continuous supply of water throughout the year and are therefore targeted for hydrochemical assessment since they are used for domestic purposes especially during dry season when most surface water and shallow wells might have dried off.

### Methods of Study

Fifteen (15) water samples were collected in thoroughly cleaned polyethylene bottles of 1.5 liter capacity. Prior to collection, the bottles were washed with distilled water and subsequently rinsed thoroughly with the sample water. The samples were collected up to top without leaving any space so as to prevent premature release of dissolved gases during the transit period. The samples were preserved with few drops of Nitric acid (HNO<sub>3</sub>). After sampling, the lids of the containers were immediately replaced to minimize contamination and escape of gases. The samples were then stored in a cooler and transported to the laboratory for analysis. All analysis was carried out at a standardized laboratory using International regulatory methods. The evaluation of water quality was in accordance with regulatory standard. The approach ensures that the samples collected were tested in accordance with agreed requirements using competent personnel as well as appropriate equipment and materials.

Field Parameters such as pH, Electrical conductivity and temperature were determined on the field due to their unstable nature. These were carried out using Multi-parameter TestrIm 35 series. Cation concentrations were measured by the Buck Scientific Model 210 VGP Atomic Absorption Spectrophotometer (AAS) while the anions were analyzed using colrimetric method. The software SPSS version 17.0 was used for statistical analyses. Graphical plots (Piper and Schoeler diagrams) were employed to unravel the hydrochemical characteristics and evolution of the spring water.

### Results and Discussion

Except for pH values that is mildly acidic in some locations, all the cations, anions and electrical conductivity are below the WHO (2004) recommended standards for drinking water. The water is slightly acidic to slightly alkaline and could be classified as fresh water [11,12]. These characteristics make the water suitable for both domestic and industrial usage.

The cations range as follows: Ca<sup>2+</sup> (1.6 – 16.0), Mg<sup>2+</sup> (0- 7.53), Na<sup>+</sup> (5.58 – 13.32), K<sup>+</sup> (0.19-20.9), while the anions are as follows: HCO<sub>3</sub><sup>-</sup> (10.0-79.0), Cl<sup>-</sup> (5.0 – 24.0), PO<sub>3</sub><sup>-</sup> (3.7 – 14.95), SO<sub>4</sub><sup>2-</sup> (4.0 – 9.2), NO<sub>3</sub><sup>-</sup> (0.12 – 9.27) (Table 1). The mean concentration of the cations is in the order Na<sup>+</sup> > Ca<sup>2+</sup> > K<sup>+</sup> > Mg<sup>2+</sup> while for anions, HCO<sub>3</sub><sup>-</sup> > Cl<sup>-</sup> > PO<sub>3</sub><sup>-</sup> > SO<sub>4</sub><sup>2-</sup> > NO<sub>3</sub><sup>-</sup>. 47% of the water are alkali water predominantly SO<sub>4</sub><sup>2-</sup> - Cl<sup>-</sup>, 20% are alkali water predominantly HCO<sub>3</sub><sup>-</sup>; 27% are earth-alkaline water predominantly SO<sub>4</sub><sup>2-</sup> while 7% are earth alkaline water predominantly HCO<sub>3</sub><sup>-</sup>; and SO<sub>4</sub><sup>2-</sup>. Table 2 is the descriptive statistical summary of the analytical data.

The water is mildly acidic to alkaline due to dissociation of bicarbonate and the following water types: Na-SO<sub>4</sub>-Cl, Na-HCO<sub>3</sub>, Ca-Na-SO<sub>4</sub> and Ca-Mg-HCO<sub>3</sub>-SO<sub>4</sub> which is a reflection of geology and

S/N	Sample ID	Temp °C	pH	Conduct µs/cm	Suspen Solid	Total Hard	Ca <sup>2+</sup>	Mg <sup>2+</sup>	Na <sup>+</sup>	K <sup>+</sup>	Cl <sup>-</sup>	PO <sub>3</sub> <sup>-</sup>	SO <sub>4</sub> <sup>-</sup>	NO <sub>3</sub> <sup>-</sup>	HCO <sub>3</sub> <sup>-</sup>
1	EPS 1	24.4	4.6	74.0	91.0	62.0	12.4	7.53	7.92	0.76	18.0	5.15	8.46	bdl	40.0
2	EPS 2	25.0	6.5	41.0	82.0	46.0	6.8	7.05	6.3	1.52	14.0	10.05	7.21	0.86	30.0
3	EPS 3	24.4	4.3	36.0	84.0	30.0	10.8	0.73	7.02	0.57	12.0	4.4	7.46	bdl	25.0
4	EPS 4	24.1	5.3	24.0	90.0	10.0	4.4	0	6.48	1.33	13.0	12.25	8.21	bdl	25.0
5	EPS 5	24.7	6.0	42.0	98.0	19.0	4.4	1.74	6.66	2.47	22.0	14.7	7.21	bdl	20.0
6	EPS 6	24.7	7.6	43.0	88.0	17.0	3.6	1.78	7.38	1.71	24.0	5.15	6.72	0.23	25.0
7	EPS 7	24.5	7.7	42.0	90.0	6.0	1.6	0.49	9.72	2.66	19.0	12.25	7.46	bdl	15.0
8	EPS 8	23.8	5.7	90.0	94.0	59.0	16	4.62	8.1	2.47	11.0	5.4	6.22	bdl	79.0
9	EPS 9	25.4	6.6	50.0	120.0	11.0	4.4	BDL	9.18	7.22	12.0	14.95	8.93	0.12	20.0
10	EPS 10	24.2	7.0	34.0	116.0	25.0	6.4	2.19	5.58	1.14	10.0	4.9	7.46	bdl	20.0
11	EPS 11	24.2	5.0	23.0	112.0	26.0	6.0	2.67	5.76	0.57	15.0	9.3	9.2	0.58	20.0
12	EPS 12	24.6	6.9	125.0	119.0	40.0	8.4	4.62	13.32	20.9	20.0	3.7	8.46	9.27	10.0
13	EPS 13	26.1	6.9	26.0	101.0	8.0	2.0	0.73	6.66	0.38	5.0	4	4.0	1.21	20.0
14	EPS 14	25.9	6.9	63.0	105.0	19.0	5.2	1.46	11.34	6.46	14.0	9.8	8.70	bdl	10.0
15	EPS 15	25.3	7.0	20.0	94.0	7.0	2.0	0.49	5.76	0.19	9.0	10.8	8.46	bdl	30.0

Table 1: Results of Water Quality Analysis in Effon Psammite Ridge Spring.

	N	Range	Minimum	Maximum	Mean		Std. Deviation	Variance
	Statistic	Statistic	Statistic	Statistic	Statistic	Std. Error	Statistic	Statistic
Temperature	15	2.3	23.8	26.1	24.7533	0.17179	0.66533	0.443
pH	15	3.4	4.3	7.7	6.2667	0.27476	1.06413	1.132
Conductivity	15	105	20	125	48.8667	7.41354	28.71253	824.41
S.Solid	15	38	82	120	98.9333	3.27395	12.67994	160.781
T.Hardness	15	56	6	62	25.6667	4.75161	18.4029	338.667
Calcium	15	14.4	1.6	16	6.2933	1.0572	4.0945	16.765
Magnesium	15	7.53	0	7.53	2.4067	0.63249	2.44964	6.001
Sodium	15	7.74	5.58	13.32	7.812	0.57455	2.22523	4.952
Potassium	15	20.71	0.19	20.9	3.3567	1.36391	5.28238	27.904
Chloride	15	19	5	24	14.5333	1.34117	5.19432	26.981
Phosphate	15	11.25	3.7	14.95	8.4533	1.03112	3.9935	15.948
Sulphate	15	5.2	4	9.2	7.6107	0.33898	1.31285	1.724
Nitrate	15	9.27	0	9.27	0.818	0.61141	2.36799	5.607
Bicarbonate	15	69	10	79	25.9333	4.28338	16.58944	275.21
Valid N (listwise)	15							

Table 2: Descriptive Statistical Summary of the Analytical Data.

climate of the study area. The mildly acidic to alkaline indices revealed that the springs water has undergone ion exchange between Na + K ions in the water with Ca and Mg of soil during the residence time of the water. Generally, the low values of the conductivity are mainly attributed to geochemical processes prevailing in the area. The springs are being recharged from recent precipitation that has low water-rock interactions and low residence time within the aquiferous zones. The hydrochemical trend signifies low mineralized water with low water-

rock interactions and residence time. Based on these water types and the presence of Na<sup>+</sup>, concentration of cations is geogenic in origin and might have come from the interaction of water and the rock or introduced from weathering of rocks into the spring water.

### Water chemistry and types

The result of chemical analysis was used to classify the water types present in the Effon Psammite Springs of the study area. Based on the predominance of both cations and anions, a plot on the Piper's Trilinear

diagram was made. Trilinear plotting systems developed by Piper [13] were used in the study of the water chemistry and quality. Piper diagrams are a combination of cation and anion triangles that lie on a common baseline. A diamond shape between them is used to replott the analyses as circles whose areas are proportional to their TDS [14-20]. The position of an analysis that is plotted on a piper diagram can be used to make a tentative conclusion as to the origin of the water represented by the analysis. The diamond part of a piper diagram may be used to characterize different water types. Generally, groundwater can be divided into four basic types according to their placement near the four corners of the diamond. Water that plots at the top of the diamond is high in  $Ca^{2+} + Mg^{2+}$  and  $Cl^{-} + SO_4^{2-}$ , which results in an area of permanent hardness. Water that plots near the left corner is rich in  $Ca^{2+} + Mg^{2+}$  and  $HCO_3^{-}$  and is the region of water of temporary hardness [20-26]. Water plotted at the lower corner of the diamond is primarily composed of alkali carbonates ( $Na^{+} + K^{+}$  and  $HCO_3^{-} + CO_3^{2-}$ ). Water lying near the right-hand side of the diamond may be considered saline ( $Na^{+} + K^{+}$  and  $Cl^{-} + SO_4^{2-}$ ). The water types in the study area were thus designated according to the area in which they occur on the diagram segments (Figure 3). These diagrams reveal the analogies, dissimilarities and different types of waters in the study area, which are shown in Figures 3 and 4. From the interpretation of this plot, several water types were delineated [27-33]. The water types are Na- $SO_4$ -Cl, Na- $HCO_3$ , Ca-Na- $SO_4$ , Ca-Mg- $HCO_3$ - $SO_4$ . Based on these water types and the presence of  $Na^{+}$ , concentration of cations is geogenic in origin and might have come from the interaction of water and the rock or introduced from weathering of rocks into the groundwater. Piper Trilinear and Durov diagrams (Figure 3 and 4 shows the water types while (Figure 5) is the Stiff diagram showing chemical ions of Effon Spring Water) [34,35].

Results also show that strong correlation exist between  $Na^{+}$  and EC




Figure 3: Piper Trilinear Diagrams for Spring Water Samples.




Figure 4: Durov Diagram of the Effon Psammite Spring Water.

	Cond.	SS	TH	pH	Ca <sup>2+</sup>	Mg <sup>2+</sup>	Na <sup>+</sup>	K <sup>+</sup>	Cl <sup>-</sup>	PO <sub>3</sub> <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>	NO <sub>3</sub> <sup>-</sup>
Cond.	1											
SS	0.28	1										
TH	0.62	-0.14	1									
pH	0.02	0.21	-0.48	1								
Ca <sup>2+</sup>	0.59	-0.11	0.95	-0.62	1							
Mg <sup>2+</sup>	0.56	-0.03	0.9	0.25	0.63	1						
Na <sup>+</sup>	0.79	0.35	0.12	0.29	0.12	0.15	1					
K <sup>+</sup>	0.79	0.56	0.13	0.27	0.09	0.21	0.86	1				
Cl <sup>-</sup>	0.34	-0.12	0.14	0.1	-0.01	0.21	0.35	0.31	1			
PO <sub>3</sub> <sup>-</sup>	0.36	0.02	-0.48	0.11	-0.48	-0.3	-0.08	-0.11	0.13	1		
SO <sub>4</sub> <sup>2-</sup>	0.12	0.31	0.04	-0.23	0.04	0.13	0.24	0.29	0.29	0.4	1	
NO <sub>3</sub> <sup>-</sup>	0.93	0.43	0.5	0.17	0.66	0.28	0.87	0.92	0.3	-0.52	0.9	1

Table 3: Correlation analysis results for the samples.

( $r=0.793$ ),  $K^{+}$  and EC ( $r=0.905$ ), TH and  $Mg^{2+}$  ( $r=0.901$ ),  $Na^{+}$  and  $K^{+}$  ( $r=0.857$ ),  $K^{+}$  and  $NO_3^{-}$  ( $r=0.909$ ),  $Ca^{2+}$  and  $Mg^{2+}$  ( $r=0.632$ ),  $Ca^{2+}$  and  $NO_3^{-}$  ( $r=0.6580$ ). Weaker correlations were obtained between  $Ca^{2+}$  and EC ( $r=0.589$ ),  $Mg^{2+}$  and EC ( $r=0.561$ ),  $K^{+}$  and Suspended solids ( $r=0.563$ ) (Table 3) [36].

### Conclusion

This investigation has revealed that except for pH values that are mildly acidic to slightly alkaline in some locations, the physico-chemical characteristics of the spring waters are below the WHO recommended standards for drinking water. The low values of the conductivity are mainly attributed to geochemical processes prevailing in the area. These characteristics make the water suitable for both domestic and industrial usage. The mean concentrations of the cations follow the order:  $Na^{+} > Ca^{2+} > K^{+} > Mg^{2+}$  while for anions,  $HCO_3^{-} > Cl^{-} > PO_3^{-} > SO_4^{2-} > NO_3^{-}$ . The water is mildly acidic to alkaline due to dissociation of bicarbonate with the following water types: Na- $SO_4$ -Cl, Na- $HCO_3$ , Ca-Na- $SO_4$  and Ca-Mg- $HCO_3$ - $SO_4$  which are a reflection of geology and climate of the area.

The mildly acidic to alkaline indices revealed that the spring water has undergone ion exchange between Na + K ions in the water with Ca and Mg of soil during the residence time of the water. The springs are being recharged from recent precipitation that has low water-rock interactions and low residence time within the aquiferous zones. The hydrochemical trend signifies low mineralized water with low water-rock interactions and residence time. Based on these water types and the presence of  $Na^{+}$ , the concentrations of cations are geogenic in origin and might have come from the interaction of water and the rock or introduced from weathering of rocks into the spring water.

It is recommended that effective development of springs should involve thorough examination of their seasonal discharges, including during the summer. In case water supply exceeds use, the surplus waters may be stored for future use in horticulture and to irrigate crop land. Since springs are yet to receive substantial attention, care must be taken to protect the spring from contamination.

### References

- Naik PK, Awasthi AK, Mohan PC (2002) Springs in a headwater basin in the Decan Trap country of the Western Ghats, India. Hydrogeology Journal 10: 553-565.
- Maxey GB (1964) Hydrogeology. In: Chow VT Handbook of Applied Hydrology. McGraw-Hill, New York.
- King M (2008) Bottled Water-Globa IndustryGuide-New Research Report on Companies and Markets.
- Aniah EJ, Eja EI, Out JE, Ushie MA (2009) Patronage of ecotourism potential

- as a strategy for sustainable tourism development in Cross River State, Nigeria. *J Geography and Geology* 1: 20-27.
5. Berner EK, Berner RA (1981) The global water cycle. Prentice Hall, Eaglewood Cliffs.
  6. Gorham E (1961) Factors influencing supply of major ions to inland waters with special references to the atmosphere. *Geol Soc Am Bull* 72: 795-840.
  7. Asiwaju-Bello YA, Akande OO (2004) Urban groundwater pollution: Case study of a disposal sites in Lagos metropolis. *Journal of water resources* 12: 22-26.
  8. Ige OO, Bale RB, Olasehinde PI (2008) Physio-chemical characteristics of water sources in Imeko, Southwestern, Nigeria. *Water res* 18: 32-36.
  9. Jones HA, Hockey RD (1964) The Geology of part of Southwestern Nigeria. *GSN Bulletin* 17: 61.
  10. Rahaman MA (1976) Review of the basement geology of Southwestern Nigeria. In *geology of Nigeria*. Elizabeth publishing Company, Nigeria.
  11. Ezeigbo HI (1989) Groundwater quality problems in parts and Sons Inc, Nigeria. *J.Min and Geology* 25: 1-4.
  12. Todd DK (1980) *Groundwater Hydrology*, (2<sup>nd</sup> Edition) John Willy and Sons Inc, New York.
  13. Piper AM (1944) A graphical procedure in the geochemical interpretation of water analysis. *American Geophysical Union Transactions* 25: 914-923.
  14. Abimbola AF, Odokoya AM, Olatunji AS (2002) Influence of bedrock on the hydrogeochemical characteristics of groundwater in Northern part of Ibadan metropolis, SW Nigeria. *Water Resources Journal* 9: 1-6.
  15. Adekunle IM, Adetunji MT, Banjoko OB (2007) Assessment of groundwater quality in a typical rural. *Int J Environ Res Public Health* 4: 307-18.
  16. Aderogba KA (2005) Ground Water Development in Nigeria: A Case Study of Abeokuta-Ewekoro-Ifo-Ota-Agbara Axis in Ogun State, Nigeria. *Int J Environ* 1: 51-68.
  17. Adeyemi GO, Adesile AO, Obayomi OM (2003) Chemical characteristics of some Well Water in Ikire, Southwestern Nigeria. *Water Resources Journal of the Nigerian Association of Hydrogeologists (NAH)* 14: 12-17.
  18. Aghazadeh N, Mogaddam NN (2010) Assessment of groundwater Quality and its suitability for Drinking and Agricultural uses in Oshnavieh Area, Northwest of Iran. *Journal of Environmental protection* 1: 30-40.
  19. Aiyegbusi MS, Aichi AU, Ugwuja OAC, Oteri AU (2010) Assessment of shallow groundwater quality in Ilesha and its environs. *Water Resources* 20: 15-28.
  20. Akujieze CN, Coker SJL, Oteze GE (2003) Groundwater in Nigeria-a millennium experience-distribution, practice, problems and solutions. *Hydrogeology* 11: 259-274.
  21. Brady NC, Weil RR (2002) The nature and properties of soils. New Jersey, Prentice-Hall.
  22. Datta PS, Tyagi SK (1996) Major ion chemistry of groundwater in Delhi area: Chemical weathering processes and groundwater flow regime. *Journal of Geological Society of India* 47: 179-188.
  23. Davies SN, DeWeist RJM (1966) *Hydrogeology*. New York: John Wiley and Sons.
  24. Drever JI (1988) *The geochemistry of natural water* (2<sup>nd</sup> Edition): Englewood Cliffs, New Jersey, prentice Hall.
  25. Freeze RA, Cherry AJ (1979) *Groundwater*. Prentice-Hall, Englewood Cliffs, NJ, 604.
  26. Hem JD (1985) Study and interpretation of the chemical characteristics of natural water. US Geological Survey Water-supply.
  27. Mayers LW (2005) *Urban Water Supply: Handbook*. New Eastern Ltd, New Delhi, India.
  28. Nagaraju A, Sureh S, Killham K, Edwards H (2006) Hydrogeochemistry of Waters of Mangampeta Barite Mining Area. Cuddapah Basin, Andhra Pradesh, India. *Turkish Journal of Engineering and Environment Sciences* 30: 203-219.
  29. Offodile ME (2002) Groundwater study and development in Nigeria. Mecon Eng Services Ltd, Jos, Nigeria.
  30. Olatunji AS, Tijani MN, Abimbola AF, Oteri AU (2001) Hydrogeochemical evaluation of water resources of Oke-Agbe Akoko, SW Nigeria. *Water Resources Journal* 12: 81-87.
  31. Raghunath IM (1987) *Groundwater*. Second edition; Wiley Eastern Ltd, New Delhi, India.
  32. Ramesh K, Elango L (2010) Groundwater quality and its suitability for domestic and agricultural use in Tondiar river basin, Tamil Nadu, India. *Environ Monit Assess* 184: 3887-3899.
  33. Richards LA (1954) *Diagnosis and improvement of Saline and Alkali Soils*. Agricultural Handbook 60, USDA and IBH publishing Co Ltd, New Delhi, India.
  34. Schlesinger WH (2004) Better living through biogeochemistry. *Ecology* 85: 2402-2407.
  35. Schoeller H (1967) *Geochemistry of groundwater*. An international guide for research and practice. UNESCO.
  36. World Health Organization (2004) *Guidelines for Drinking Water Quality: 3<sup>rd</sup> Incorporation of 1<sup>st</sup> and 2<sup>nd</sup> Addenda*, Geneva.

**Citation:** Nwankwoala HO, Nwaogu C, Bolaji TA, Uzoegbu MU, Abrakasa S, et al., (2015) Geochemistry and Quality Characterization of Effon Psammite Ridge Spring Water, Southwestern Nigeria. *J Geol Geophys* 4: 229. doi:[10.4172/jgg.1000229](https://doi.org/10.4172/jgg.1000229)

### OMICS International: Publication Benefits & Features

#### Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

#### Special features:

- 700 Open Access Journals
- 50,000 Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus, Google Scholar etc.
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsgroup.org/journals/submission>